

2363eP - Treatment intensification in real-world metastatic hormone-sensitive prostate cancer: contemporary patterns across four countries

Mario Uccello¹, Christine Mai², Hilary Worton³, Stergios Boussios⁴, Bertrand de Buhren², Malgorzata Gozdzik², Maura De Almeida²
¹Royal United Hospitals Bath NHS Foundation Trust, Bath, UK; ²APLUSA, Lyon, France; ³Aplusa UK, London, UK; ⁴UOI - University of Ioannina, Ioannina, Greece

BACKGROUND



Treatment intensification beyond androgen deprivation therapy (ADT) is now widely recommended for patients with metastatic hormone-sensitive prostate cancer (mHSPC). However, the adoption of intensified regimens in routine clinical practice remains heterogeneous. We evaluated contemporary first-line treatment patterns across four countries, with particular focus on differences according to metastatic presentation and geographic setting.

METHODS

Anonymised clinical charts of patients with mHSPC treated between 2025 and 2026 were collected from oncologists in China, France, Germany and Japan. First-line systemic therapy was classified as ADT alone, androgen receptor pathway inhibitor (ARPI) + ADT, taxane + ADT, taxane + ARPI + ADT, or other. Treatment patterns were assessed overall and according to country and de novo versus recurrent metastatic presentation, with differences evaluated using chi-square tests.

Uccello M. declares no conflict of interest.
Contact: mario_uccello@hotmail.it

RESULTS



CONCLUSION

- Treatment intensification was used in most patients with mHSPC.
- ARPI + ADT was the most frequently used first-line regimen.
- Triplet therapy was more commonly used in de novo than in recurrent metastatic disease.
- Treatment patterns varied significantly across countries, indicating persistent geographic differences in real-world practice.